

For any questions or concerns, call 781-383-8380.

For fever greater than 100.4°F in an infant < 3 months old, contact your pediatrician first.

Do not alternate acetaminophen and ibuprofen.



### Acetaminophen (generic Tylenol)

May be given every 4 hours as needed. Do not exceed 5 doses in a 24 hour period.

Child's weight (pounds)	6-11	12-17	18-23	24-35	36-47	48-59	60-71	72-95	96+
Syrup: 160 mg/5ml	1.25ml	2.5ml	3.75ml	5ml (1tsp)	7.5ml	10ml	12.5ml	15ml	20ml
Chewable: 160 mg tablets				1 tabs	1.5 tabs	2 tabs	2.5 tabs	3 tabs	4 tabs
Adult: 325 mg tablets						1 tab	1 tab	1.5 tab	2 tabs
Adult: 500mg tablets								1 tab	1 tab

### Ibuprofen (generic Motrin)

May be given every 6-8 hours as needed. Do not use under 6 months of age.

Child's weight (pounds)	12-17	18-23	24-35	36-47	48-59	60-71	72-95	96+
Infant Drops: 50mg/1.25ml	1.25ml	1.875ml	2.5ml	3.75ml	5 ml			
Liquid: 100mg/5ml	2.5ml	4ml	5ml	7.5ml	10ml	12.5ml	15ml	20ml
Chewable: 100mg tabs			1 tab	1.5 tab	2 tab	2.5 tab	3 tab	4 tab
Junior strength: 100mg tabs					2 tab	2.5 tab	3 tab	4 tab
Adult: 200mg tabs					1 tab	1 tab	1.5 tab	2 tab

### Benadryl (Diphenhydramine)

May be given every 6-8 hours as needed. Do not use under 1 year of age.

Child's weight (pounds)	20-24	25-47	38-49	50-99	100+
Liquid: 12.5/5ml	4ml	5ml	7.5ml	10ml	
Chewable: 12.5mg		1 tab	1.5 tab	2tab	4 tab
Tablets: 25mg		0.5 tab	0.5 tab	1 tab	2 tab
Capsules: 25/mg				1 cap	2 cap